

DOWNER -- NEW JERSEY STATE SOIL



Downer Soil Profile

Surface layer: dark grayish brown loamy sand
Subsurface layer: grayish brown sandy loam
Subsoil - upper: yellowish brown gravelly sandy loam
Subsoil - lower: yellowish brown sand and coarse sand

Downer soils are used mostly as woodland. The natural vegetation consists of mixed oaks, hickory, and scattered pines. Some areas are cultivated for high-value vegetable and fruit crops that are usually irrigated. Downer soils occur on 291,319 acres in New Jersey. These soils are dominantly in the 11 counties of southern New Jersey.

Downer soils are on uplands and formed in sandy, highly quartzose Northern Atlantic Coastal Plain deposits. They have medium natural fertility and require fertilization for maximum crop production. The Downer series was established in Gloucester County, New Jersey, in 1960.



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD).

 **NRCS** Natural Resources Conservation Service
Helping People Help the Land